

# WHAT'S ON THE MENU?

## DECIDING WHAT IS AVAILABLE TO THE GROUP\*

David S. Ahn<sup>†</sup>  
University of California, Berkeley

Christopher P. Chambers<sup>‡</sup>  
California Institute of Technology

February 2008

### Abstract

We consider the problem of aggregating menus. Each agent submits a menu of options, which the social rule aggregates into a single menu. First, we study a general environment where menus are arbitrary subsets, and characterize a union rule which takes the union of individual menus. Second, we study a probabilistic environment where menus are closed convex subsets of lotteries over options, and characterize a rule which takes a convex combination over individual menus with fixed weights.

## 1 Introduction

The basic problem of deciding a common set of options for a number of agents arises frequently. Consider the following:

- Two friends want to meet at a restaurant for dinner. One is on a diet and hopes to avoid unhealthy options, while the other has just received a promotion and hopes to celebrate by indulging in a rich meal.
- A government designs a mandatory pension system, to be funded through implicit forced savings taxed on current income. Some residents prefer the flexibility for current consumption afforded with smaller rates of forced savings. Others prefer larger rates to moderate their temptation to overconsume in the present. A single common system must be incorporated into the tax code.

---

\*The authors thank Federico Echenique, Ivana Komunjer, Rolf Schneider, and seminar participants at Carlo Alberto and CORE for helpful comments. We are especially grateful to Todd Sarver for detailed comments and suggestions.

<sup>†</sup>Department of Economics, University of California, 549 Evans Hall #3880, Berkeley, CA 94720-3880. Email: [dahn@econ.berkeley.edu](mailto:dahn@econ.berkeley.edu).

<sup>‡</sup>Division of Humanities and Social Sciences, 228-77, California Institute of Technology, Pasadena, CA 91125. Email: [chambers@hss.caltech.edu](mailto:chambers@hss.caltech.edu).

- Many laws are basic restrictions on behavior applied consistently to everyone. Citizens disagree on limits of abortion rights, the right to free speech, and so on, but a single set of restrictions must be reached.
- Nations often meet to ratify treaties to limit their behavior. For example, the Geneva Conventions dictate rules of engagement during war. Some countries prefer to restrict torture of prisoners and attacks on civilians, while others would like more liberal rules allowing these practices.

Kreps (1979) introduced an axiomatic theory of decision making for a single agent over menus, which is extended by Dekel, Lipman, and Rustichini (2001) and Gul and Pesendorfer (2001). These theories incorporate a particular agent’s desire for larger menus to provide flexibility or for smaller menus to provide commitment. A common theme in the prior examples is a social tension between flexibility and commitment across many agents. We study rules for resolving this tension.

In particular, we focus attention to a simple setting where each agent proposes or submits a menu, and the social rule aggregates these submissions into a common menu. An alternative approach might aggregate entire preferences over menus. For example, a variant of the classical Harsanyi (1955) Aggregation Theorem, e.g. De Meyer and Mongin (1995), implies that a Paretian, independent, and monotone social aggregation of independent and monotone preferences over sets of lotteries can be represented with a convex combination of the subjective state spaces and measures which represent individuals’ preferences. One difficulty with implementing this rule is obtaining each individual’s preference over menus. The main substantive motivation for the current setting is its relative simplicity. Each agent submits a single menu, rather than a rank order of the entire power set of menus.

In this paper, we consider two structural environments. First is the general environment where no structure is imposed on the space of outcomes, which is analogous to Kreps (1979). Here, we characterize a rule which returns the union of the individually proposed menus. Second is a probabilistic environment where the outcome space is the space of lotteries over deterministic outcomes, which is analogous to Dekel, Lipman, and Rustichini (2001) and Gul and Pesendorfer (2001) and provides some useful linear structure. Here, we characterize a rule which returns a convex combination with fixed weights over the individually proposed menus.

While this paper is motivated by temptation and flexibility, there are other interpretations of preferences over subsets, e.g. in the literatures on voting or matching. Barberà, Bossert, and Pattanaik (2004) provide a good review of these interpretations. The interpretation and appropriateness of the following aggregation axioms are sensitive to how subsets are interpreted, so some care should be exercised before applying the results to alternative settings.

For example, the general model considered here is formally similar to approval voting (Brams and Fishburn 1978, Weber 1995), where voters submit ballots of acceptable candidates from each voter and outputs as winners the candidates which appear on the most ballots. Aside from the

obvious difference in interpretation, the resulting representation is also distinct.<sup>1</sup> The union rule considered in this paper would violate the Consistency condition required to characterize approval voting across populations (Fishburn 1978), while approval voting would violate the Monotonicity condition used here.

A final interpretation of the environment is a resolution of competing opinions or conjectures. The general model can be viewed in terms of a principal asking various advisors about what they view as the possible states of the world, while the probabilistic model corresponds to asking advisors for their sets of beliefs over states. A corollary contribution of the paper is to provide characterizations of different belief aggregators in these settings.

## 2 General model

We begin by describing the general model. Let  $N = \{1, \dots, n\}$  denote a finite set of agents and  $X$  denote an arbitrary set of alternatives. In general  $\mathcal{X} \subseteq 2^X$  denotes some universe of admissible subsets or menus. In this section, we will take  $\mathcal{X} = 2^X \setminus \{\emptyset\}$  to be the family of all nonempty subsets of  $X$ . A rule is a function  $F : \mathcal{X}^N \rightarrow \mathcal{X}$  assigning a menu to every profile of menus. One such rule is the **union rule** which maps each profile to the union of its components,  $F(A_1, \dots, A_n) = \bigcup_{i \in N} A_i$ . This is a liberal rule in the following sense: if any agent desires the freedom to choose a particular option, the social menu will include that option.

We now consider some restrictions on rules.

**Axiom 1** (Unanimity). For all  $A \in \mathcal{X}$ ,  $F(A, \dots, A) = A$ .

Unanimity is a minimal efficiency criterion. If every individual agrees on the same menu, then that menu should be the social outcome.

**Axiom 2** (Anonymity). For any bijection  $\tau : N \rightarrow N$  and profile  $(A_1, \dots, A_n) \in \mathcal{X}^N$ ,  $F(A_1, \dots, A_n) = F(A_{\tau(1)}, \dots, A_{\tau(n)})$ .

This classical condition, which makes the identities of the agents irrelevant, captures some notion of equity. As the proofs will make clear, it is also very technically useful in this setting.

**Axiom 3** (Monotonicity). If  $A_i \subseteq B_i$  for all  $i \in N$ , then  $F(A) \subseteq F(B)$ .

Monotonicity asserts that if every agent submits a larger menu with more flexibility (or a smaller one with more commitment), then the social outcome should also allow for more flexibility (or impose more commitment). This condition is directly related to the interpretation of menus through commitment and flexibility.

**Axiom 4** (Disjoint Additivity). If  $A_i \cap B_i = \emptyset$  for all  $i \in N$ , then  $F(A_i \cup B_i, \dots, A_n \cup B_n) = F(A_1, \dots, B_n) \cup F(B_1, \dots, B_n)$ .

---

<sup>1</sup>Moreover, approval voting is generally characterized with conditions applied across populations, while we focus on a single population setting.

We can also consider the following stronger condition, applied at the level of the individual submission rather than the entire profile: if  $A_i \cap B_i = \emptyset$  for some  $i \in N$ , then

$$F(A_i \cup B_i, (C_j)_{j \neq i}) = F(A_i, (C_j)_{j \neq i}) \cup F(B_i, (C_j)_{j \neq i}).$$

Thusly stated, Disjoint Additivity resembles a path independence or separability condition. The social menu induced by a profile of individual menus can be decomposed and analyzed through disjoint individual submenus. One way to understand this condition is suppose we split the domain  $X = X' \cup X''$ , with  $X', X''$  disjoint, then ask some agent  $i$  to submit a menu  $A_i \subseteq X'$  and  $B_i \subseteq X''$  from each subdomain. For example, these menus might respectively be the stocks or bonds she would like to have as investments for her retirement account, conditional on only one kind of asset being available. If she selects a conditional menu  $A_i \subseteq X'$  of stocks and a conditional menu  $B_i \subseteq X''$  of bonds, then the social menu that would have been offered if she had submitted the union  $A_i \cup B_i$  of stock and bond menus is exactly the union of the social menus that are offered based on her conditional menus, fixing all other agents' submissions. Of course, someone who submits  $A_i$  from  $X'$  and  $B_i$  from  $X''$  might submit a separate menu not equal to  $A_i \cup B_i$  from the grand domain  $X$ : for example, she finds some option  $a \in A_i$  tempting against  $B_i$  in a way that it is not against other choices in  $A_i$ . The axiom asserts only that if she were to submit the union  $A_i \cup B_i$  from the grand domain, then the resulting social menu would take the union of the conditional social menus.

**Theorem 1.** *Suppose  $|X| \geq |N|$ . A rule  $F : \mathcal{X}^N \rightarrow \mathcal{X}$  satisfies Unanimity, Anonymity, Monotonicity, and Disjoint Additivity if and only if  $F(A_1, \dots, A_n) = \bigcup_{i \in N} A_i$ .*

*Proof.* We omit the straightforward verification of necessity, moving directly to sufficiency.

We first prove the claim for profiles of singletons. Let  $x_i \in X$  for each  $i \in N$  and  $A = \bigcup_{i \in N} \{x_i\}$ . Let  $m = |A|$  denote the number of distinct elements in the profile  $(\{x_1\}, \dots, \{x_n\})$ . We need to show  $F(\{x_1\}, \dots, \{x_n\}) = A$ . Since there are more objects than agents,  $|X| \geq |N|$ , there exists some residual set  $B = \{b_1, \dots, b_{n-m}\} \subseteq X \setminus \{x_1, \dots, x_m\}$  such that  $|B| = n - m$ .

One set containment,  $F(\{x_1\}, \dots, \{x_n\}) \subseteq F(A, \dots, A) \subseteq A$ , follows from Monotonicity and Unanimity. This also implies that  $F(\{x_1\}, \dots, \{x_n\})$  is disjoint from  $B$ . So, to prove that  $A \subseteq F(\{x_1\}, \dots, \{x_n\})$ , it suffices to show that  $A \cup B \subseteq F(\{x_1\}, \dots, \{x_n\}) \cup B$ .

To this end, we may assume, by reordering and appealing to Anonymity, that the first  $m$  components of the profile are distinct:  $x_i \neq x_j$  for all  $i, j \leq m$ . Consider the profile

$$(\{y_1\}, \dots, \{y_n\}) = (\{x_1\}, \{x_2\}, \dots, \{x_m\}, \{b_1\}, \{b_2\}, \dots, \{b_{n-m}\}).$$

By construction,  $y_i \neq y_j$  whenever  $i \neq j$  and  $\{y_1, \dots, y_n\} = A \cup B$ . Let  $\tau^i$  denote the  $i$ -th cyclic

permutation on  $N$ .<sup>2</sup>

$$\begin{aligned}
A \cup B &= F(A \cup B, \dots, A \cup B), && \text{by Unanimity} \\
&= F\left(\bigcup_{i \in N} \{y_{\tau^i(1)}\}, \dots, \bigcup_{i \in N} \{y_{\tau^i(n)}\}\right) \\
&= \bigcup_{i \in N} F(\{y_{\tau^i(1)}\}, \dots, \{y_{\tau^i(n)}\}), && \text{by } n \text{ applications of Disjoint Additivity} \\
&= F(\{y_1\}, \dots, \{y_n\}), && \text{by Anonymity} \\
&\subseteq F(\{x_1\} \cup B, \dots, \{x_n\} \cup B), && \text{by Monotonicity} \\
&= F(\{x_1\}, \dots, \{x_n\}) \cup F(B, \dots, B), && \text{by Disjoint Additivity} \\
&= F(\{x_1\}, \dots, \{x_n\}) \cup B, && \text{by Unanimity.}
\end{aligned}$$

Moving to the general case, take an arbitrary profile  $(A_1, \dots, A_n)$ . By Monotonicity and Unanimity,  $F(A_1, \dots, A_n) \subseteq F(\bigcup_{i \in N} A_i, \dots, \bigcup_{i \in N} A_i) = \bigcup_{i \in N} A_i$ . For the other direction, suppose  $x \in \bigcup_{i \in N} A_i$ . Then  $x \in A_i$  for some  $A_i$ , so there exists a selection  $(x_1, \dots, x_n) \in \prod_{i \in N} A_i$  such that  $x \in \{x_1, \dots, x_n\}$ . But  $x \in \{x_1, \dots, x_n\} = F(\{x_1\}, \dots, \{x_n\}) \subseteq F(A_1, \dots, A_n)$ , by the previous step and Monotonicity. So, we have  $\bigcup_{i \in N} A_i \subseteq F(A_1, \dots, A_n)$ .  $\square$

The simple verification of the logical independence of the axioms is omitted.

The hypothesis that the number of objects is at least as large as the number of agents is indispensable.

**Example 1.** Let  $X = \{x, y\}$  and  $N = \{1, 2, 3\}$ . If there exists some  $i \in N$  such that  $A_i = \{x, y\}$ , let  $F(A_1, A_2, A_3) = \{x, y\}$ . There are four remaining profiles to consider, up to permutation of agents:

$$\begin{aligned}
F(\{x\}, \{x\}, \{x\}) &= \{x\} \\
F(\{x\}, \{x\}, \{y\}) &= \{x\} \\
F(\{x\}, \{y\}, \{y\}) &= \{y\} \\
F(\{y\}, \{y\}, \{y\}) &= \{y\}.
\end{aligned}$$

This example satisfies Unanimity, Anonymity, Monotonicity, and Disjoint Additivity.

The theorem can also be proved with a stronger Additivity condition whose hypothesis does not require that announced menus be disjoint, in which case the cardinality hypothesis is no longer required. Such an axiom implies both Disjoint Additivity and Monotonicity.

---

<sup>2</sup>So  $\tau^i(k) = [k + i] \bmod n$ .

### 3 Probabilistic model

In this section, suppose  $X$  is a finite set. Let  $\Delta X$  denote the set of lotteries over  $X$ . We now take  $\mathcal{X} = \mathcal{K}(\Delta X)$ , the nonempty, closed, and convex subsets of  $\Delta X$ .<sup>3</sup> Define the convex combination of two sets  $A, B \in \mathcal{K}$  with weight  $\alpha \in [0, 1]$  by

$$\alpha A + (1 - \alpha)B = \{\alpha p + (1 - \alpha)q : p \in A, q \in B\}.$$

The additional stochastic structure here affords new kinds of aggregation rules. For example, the **weighting rule** takes a convex combination, with fixed weights across agents, of the individual menus:  $F(A_1, \dots, A_n) = \sum_{i \in N} \mu_i A_i$ , for some  $\mu \in \Delta N$ . This rule exploits objective randomization in a manner similar to how randomized allocation of an indivisible object improves ex ante equity.

The weighting rule fails Disjoint Additivity, but satisfies the following condition, which invokes the linear structure of the space of lotteries.

**Axiom 5** (Mixture Linearity). For all  $(A_1, \dots, A_n), (B_1, \dots, B_n) \in \mathcal{X}^N$  and  $\alpha \in [0, 1]$ ,

$$F(\alpha A_1 + (1 - \alpha)B_1, \dots, \alpha A_n + (1 - \alpha)B_n) = \alpha F(A_1, \dots, A_n) + (1 - \alpha)F(B_1, \dots, B_n).$$

This axiom is an analog of independence for individual choice over menus of lotteries, as used Dekel, Lipman, and Rustichini (2001). Its normative appeal is also analogous, in terms of the timing of resolution of uncertainty. In this interpretation, the combination  $(\alpha A_1 + (1 - \alpha)B_1, \dots, \alpha A_n + (1 - \alpha)B_n)$  is viewed as a randomization, say generated by an  $\alpha$ -weighted coin, between the profiles  $(A_1, \dots, A_n)$  and  $(B_1, \dots, B_n)$ . Then the social menu over these randomized submissions is the randomization of the social menu induced by the submissions  $(A_1, \dots, A_n)$  and  $(B_1, \dots, B_n)$ . So, a Mixture Linear rule is invariant to whether the coin is flipped before or after the point of aggregation.

Replacing Disjoint Additivity with Mixture Linearity, we characterize the weighting rule with a uniform weighting over agents.

**Theorem 2.** *A rule  $F : \mathcal{X}^N \rightarrow \mathcal{X}$  satisfies Unanimity, Anonymity, and Mixture Linearity if and only if  $F(A_1, \dots, A_n) = \sum_{i \in N} \frac{1}{n} A_i$ .*

*Proof.* Let  $(A_1, \dots, A_n) \in \mathcal{X}^N$ . Recall  $\tau^i : N \rightarrow N$  denoted the  $i$ -th cyclic permutation on  $N$ .

---

<sup>3</sup>We could also allow for sets which are not convex by assuming that  $F(A_1, \dots, A_n) = F(\text{conv}(A_1), \dots, \text{conv}(A_n))$ , where  $\text{conv}(A_i)$  denotes the convex hull of  $A_i$ .

Then:

$$\begin{aligned}
F(A_1, \dots, A_n) &= F(A_{\tau^i(1)}, \dots, A_{\tau^i(n)}), && \text{by Anonymity} \\
&= \sum_{i \in N} \frac{1}{n} F(A_{\tau^i(1)}, \dots, A_{\tau^i(n)}) \\
&= F\left(\frac{1}{n} \sum_{i \in N} A_{\tau^i(1)}, \dots, \frac{1}{n} \sum_{i \in N} A_{\tau^i(n)}\right), && \text{by Mixture Linearity} \\
&= F\left(\frac{1}{n} \sum_{i \in N} A_i, \dots, \frac{1}{n} \sum_{i \in N} A_i\right) \\
&= \frac{1}{n} \sum_{i \in N} A_i, && \text{by Unanimity.} \quad \square
\end{aligned}$$

Again, it is simple to verify that the three axioms are independent. The characterization makes no use of the Monotonicity condition which concerns commitment versus flexibility, and as such requires no appeal to that interpretation over others.

If the Anonymity axiom is dispensed, the weighting becomes arbitrary over agents. This is not a simple corollary of the Mixture Space Theorem or the standard Harsanyi Aggregation Theorem. In particular, the space of convex sets is somewhat intractable, and the proof moves the analysis to the space of support functions, following Dekel, Lipman, and Rustichini (2001). The consequent result can be interpreted as a version of the Aggregation Theorem over the infinite-dimensional space of sets.

**Theorem 3.** *A rule  $F : \mathcal{X}^N \rightarrow \mathcal{X}$  satisfies Unanimity, Monotonicity, and Mixture Linearity if and only if there exists  $\mu \in \Delta N$  such that  $F(A_1, \dots, A_n) = \sum_{i \in N} \mu_i A_i$ .*

*Proof.* We omit the straightforward verification of necessity and proceed to show sufficiency.

Let  $m = |X|$  and  $D = \{x \in \mathbb{R}^m : \|x\| = 1 \text{ and } \sum_i x_i = 0\}$ . For any  $A \in \mathcal{X}$ , define the support function  $\sigma_A : D \rightarrow \mathbb{R}$  of a set  $A \in \mathcal{X}$  by  $\sigma_A(s) = \max_{a \in A} a \cdot s$ . Let  $\mathcal{X}^* = \{\sigma_A : A \in \mathcal{X}\} \subseteq \mathbb{R}^D$  denote the family of all support functions induced by some set in  $\mathcal{X}$ . Define the dual function  $F^* : [\mathcal{X}^*]^N \rightarrow \mathcal{X}^*$  by

$$F^*(\sigma_{A_1}, \dots, \sigma_{A_n}) = \sigma_{F(A_1, \dots, A_n)}.$$

It suffices to prove that  $F^*(\sigma_{A_1}, \dots, \sigma_{A_n}) = \sum_{i \in N} \mu_i \sigma_{A_i}$ .

The dual function inherits mixture linearity:

$$\begin{aligned}
& F^*(\alpha\sigma_{A_1} + (1-\alpha)\sigma_{B_1}, \dots, \alpha\sigma_{A_n} + (1-\alpha)\sigma_{B_n}) \\
&= F^*(\sigma_{\alpha A_1 + (1-\alpha)B_1}, \dots, \sigma_{\alpha A_n + (1-\alpha)B_n}) \\
&= \sigma_{F(\alpha A_1 + (1-\alpha)B_1, \dots, \alpha A_n + (1-\alpha)B_n)} \\
&= \sigma_{\alpha F(A_1, \dots, A_n) + (1-\alpha)F(B_1, \dots, B_n)} \\
&= \alpha\sigma_{F(A_1, \dots, A_n)} + (1-\alpha)\sigma_{F(B_1, \dots, B_n)} \\
&= \alpha F^*(\sigma_{A_1}, \dots, \sigma_{A_n}) + (1-\alpha) F^*(\sigma_{B_1}, \dots, \sigma_{B_n}).
\end{aligned}$$

The support function also carries a form of monotonicity to the dual, since  $\sigma_A \leq \sigma_{A'}$  if and only if  $A \subseteq A'$ . So, if  $\sigma_{A_i} \leq \sigma_{B_i}$ , i.e.  $A_i \subseteq B_i$ , for all  $i \in N$ , then:

$$F^*(\sigma_{A_1}, \dots, \sigma_{A_n}) = \sigma_{F(A_1, \dots, A_n)} \leq \sigma_{F(B_1, \dots, B_n)} = F^*(\sigma_{B_1}, \dots, \sigma_{B_n}).$$

Now, for any  $s \in D$ , consider the real-valued function  $F_s^* : [\mathcal{X}^*]^N \rightarrow \mathbb{R}$  defined by

$$F_s^*(\sigma_{A_1}, \dots, \sigma_{A_n}) = [F^*(\sigma_{A_1}, \dots, \sigma_{A_n})](s).$$

This function is also mixture linear and monotonic. Note that

$$F_s^*(0, \dots, 0) = 0.$$

To see this, let  $\ell^0 = (\frac{1}{m}, \dots, \frac{1}{m})$  denote the uniform lottery over  $X$  and observe:

$$F_s^*(0, \dots, 0) = F^*(\sigma_{\{\ell_0\}}, \dots, \sigma_{\{\ell_0\}}) = \sigma_{\{\ell_0\}} = 0.$$

The argument for Theorems 1 and 2 of Dekel, Lipman, Rustichini, and Sarver (2007) can be replicated, with some obvious adjustments to account for the product space of the domain, to produce an increasing continuous linear extension  $W_s : [C(D)]^N \rightarrow \mathbb{R}$  of  $F_s^*$  to  $[C(D)]^N$ , the space of all profiles of continuous real-valued functions on  $D$ .

Therefore, by the Riesz Representation Theorem, there exists a profile  $(\mu_1^s, \dots, \mu_n^s)$  of finite nonnegative measures on  $D$  such that

$$F_s^*(\sigma_{A_1}, \dots, \sigma_{A_n}) = W_s(\sigma_{A_1}, \dots, \sigma_{A_n}) = \sum_{i \in N} \int_D \sigma_{A_i}(t) d\mu_i^s(t).$$

The measure  $\mu_i^s$  is concentrated on  $\{s\}$ . To see this, recall that  $\ell^0$  is the uniform distribution and consider the closed convex set  $A = \{\ell \in \Delta X : \ell \cdot s \leq 0\}$ . Let  $t \in D \setminus \{s\}$ . Note that there exists some  $\ell \in A$  for which  $\ell \cdot t > 0$ , so that  $\sigma_A(t) > 0$ .



Then, by Unanimity:

$$\begin{aligned}
\sigma_A(s) &= \sigma_{\ell^0}(s) \\
F_s^*(\sigma_A, \dots, \sigma_A) &= F_s^*(\sigma_{\{\ell^0\}}, \dots, \sigma_{\{\ell^0\}}) \\
\sum_{i \in N} \int_D \sigma_A(t) d\mu_i^s(t) &= \sum_{i \in N} \int_D \sigma_{\{\ell^0\}}(t) d\mu_i^s(t) \\
\sum_{i \in N} \left[ \sigma_A(t) \mu_i^s(\{s\}) + \int_{D \setminus \{s\}} \sigma_A(s) d\mu_i^s(t) \right] &= \sum_{i \in N} \left[ \sigma_{\{\ell^0\}}(s) \mu_i^s(\{s\}) + \int_{D \setminus \{s\}} \sigma_{\{\ell^0\}}(t) d\mu_i^s(t) \right] \\
\sum_{i \in N} \int_{D \setminus \{s\}} \sigma_A(t) d\mu_i^s(t) &= \sum_{i \in N} \int_{D \setminus \{s\}} \sigma_{\{\ell^0\}}(t) d\mu_i^s(t)
\end{aligned}$$

But, since  $\sigma_A > \sigma_{\{\ell^0\}}$  on  $D \setminus \{s\}$  and every  $\mu_i^s$  is nonnegative, this suffices to show  $\mu_i^s(D \setminus \{s\}) = 0$ , as the right hand side of the final equality is zero. So, let  $\mu_i^s = \mu_i^s(\{s\})$  and note that Unanimity implies  $\sum_{i \in N} \mu_i^s = 1$ .

We lastly show that  $\mu^s = \mu^t$  for all  $s, t \in D$ . As a first step, observe that if  $(\{x_1\}, \dots, \{x_n\})$  is a profile of singletons, then  $F(\{x_1\}, \dots, \{x_n\})$  is a singleton. To see this, note that by Mixture Linearity and Unanimity,

$$\sum_{i \in N} \frac{1}{n} F(\{x_{\sigma^i(1)}\}, \dots, \{x_{\sigma^i(n)}\}) = \left\{ \sum_{i \in N} \frac{1}{n} x_{\sigma^i(n)} \right\}.$$

Since the right hand side is a singleton, each component of the sum on the left hand side must also be a singleton.

To establish that  $\mu^s = \mu^t$  for all  $s, t$ , we will consider two cases, the case in which  $s = -t$  and the case in which  $s \neq -t$ . We will show that for any  $i \in N$ ,  $\mu_s^i = \mu_t^i$ .

Recall that  $\ell^0$  is the uniform distribution over  $X$ . Let  $i \in N$ . Let  $(A_1, \dots, A_n)$  be a profile of singletons such that for all  $j \neq i$ ,  $A_j = \{\ell^0\}$ , and  $A_i$  is any singleton. We claim that there exists some  $\mu^i \in [0, 1]$  such that  $F(A_1, \dots, A_n) = \mu^i A_i + (1 - \mu^i)\{\ell^0\}$ . But this follows trivially by Unanimity, Monotonicity, and the prior observation that singleton profiles map to singletons.

Now, suppose that  $s = -t$ . Let us choose  $\varepsilon$  small enough so that  $\ell^0 + \varepsilon s \in \Delta X$ . Consider the profile where  $A_i = \{\ell^0 + \varepsilon s\}$  and for all  $j \neq i$ ,  $A_j = \{\ell^0\}$ . Then  $F(A_1, \dots, A_n) = \mu^i \{\ell^0 + \varepsilon s\} + (1 - \mu^i)\{\ell^0\}$  for some  $\mu^i$ . In particular, as  $\sigma_{\ell^0}(s') = 0$ , we may conclude that  $F_s^*(\sigma_{A_1}, \dots, \sigma_{A_n}) = \mu^i \varepsilon$  and  $F_t^*(\sigma_{A_1}, \dots, \sigma_{A_n}) = -\mu^i \varepsilon$ . Consequently,  $F_s^*(\sigma_{A_1}, \dots, \sigma_{A_n}) = -F_t^*(\sigma_{A_1}, \dots, \sigma_{A_n})$ . Now,

$$\mu_i^s = \frac{\mu_i^s \varepsilon}{\varepsilon} = \frac{\mu_i^s \sigma_{A_i}(s) + \sum_{j \neq i} \mu_j^s \sigma_{A_j}(s)}{\varepsilon} = \frac{F_s^*(\sigma_{A_1}, \dots, \sigma_{A_n})}{\varepsilon}$$

and

$$\mu_i^t = \frac{\mu_i^t(-\varepsilon)}{-\varepsilon} = \frac{\mu_i^t \sigma_{A_i}(t) + \sum_{j \neq i} \mu_j^t \sigma_{A_j}(t)}{-\varepsilon} = \frac{F_t^*(\sigma_{A_1}, \dots, \sigma_{A_n})}{-\varepsilon},$$

from which we conclude  $\mu_i^t = \mu_i^s$ .

Next consider the case where  $s \neq -t$ . For sufficiently small  $\varepsilon > 0$ ,  $\ell^0 + \varepsilon(s+t) \in \Delta X$ . Then:

$$\sigma_{\{\ell^0 + \varepsilon(s+t)\}}(s) = \sigma_{\{\ell^0\}}(s) + \sigma_{\{\varepsilon(s+t)\}}(s) = \varepsilon(1+t \cdot s) = \sigma_{\{\ell^0 + \varepsilon(s+t)\}}(t)$$

Consider the profile  $(A_1, \dots, A_n)$  where  $A_i = \{\ell^0 + \varepsilon(s+t)\}$  and  $A_j = \{\ell^0\}$  for all  $j \neq i$ . As previously, let us write  $F(A_1, \dots, A_n) = \mu_i \{\ell^0 + \varepsilon(s+t)\} + (1 - \mu_i) \{\ell^0\}$ . Therefore,  $F_s^*(\sigma_{A_1}, \dots, \sigma_{A_n}) = \mu_i \varepsilon(1+t \cdot s) = F_t^*(\sigma_{A_1}, \dots, \sigma_{A_n})$ .

Then:

$$\mu_i^s = \frac{\mu_i^s \varepsilon(1+t \cdot s)}{\varepsilon(1+t \cdot s)} = \frac{\mu_i^s \sigma_{A_i}(s) + \sum_{j \neq i} \mu_j^s \sigma_{A_j}(s)}{\varepsilon(1+t \cdot s)} = \frac{F_s^*(\sigma_{A_1}, \dots, \sigma_{A_n})}{\varepsilon(1+t \cdot s)}$$

and

$$\mu_i^t = \frac{\mu_i^t \varepsilon(1+t \cdot s)}{\varepsilon(1+t \cdot s)} = \frac{\mu_i^t \sigma_{A_i}(t) + \sum_{j \neq i} \mu_j^t \sigma_{A_j}(t)}{\varepsilon(1+t \cdot s)} = \frac{F_t^*(\sigma_{A_1}, \dots, \sigma_{A_n})}{\varepsilon(1+t \cdot s)}$$

As  $F_s^*(\sigma_{A_1}, \dots, \sigma_{A_n}) = F_t^*(\sigma_{A_1}, \dots, \sigma_{A_n})$ , we may conclude  $\mu_i^s = \mu_i^t$ .

Hence, there exists a  $\mu \in \Delta N$  such that  $[F^*(\sigma_{A_1}, \dots, \sigma_{A_n})](s) = \sum_{i \in N} \mu_i \sigma_{A_i}(s)$  for all  $s \in D$ , as desired.  $\square$

Finally, we note a question our analysis leaves open, in the hope that it is resolved by future research. Monotonicity is crucial in the proof to ensure that the linear extension of the dual function is continuous, hence submits to the Reisz Representation Theorem. We are unsure whether Monotonicity is redundant: we could not prove the result without it, nor could we construct an example which satisfies the other conditions but fails Monotonicity. The subtlety of the problem will be familiar to decision theorists working with similar mathematical tools. While Monotonicity seems normatively innocuous, it would nonetheless be technically satisfying to resolve this issue conclusively.

## References

- BARBERÀ, S., W. BOSSERT, AND P. K. PATTANAIK (2004): "Ranking Sets of Objects," in *Handbook of Utility Theory: Volume 2: Extensions*, ed. by S. Barberà, P. J. Hammond, and C. Seidl, pp. 893–977, Dordrecht. Kluwer.
- BRAMS, S. J., AND P. C. FISHBURN (1978): "Approval Voting," *American Political Science Review*, 72, 831–847.
- DE MEYER, B., AND P. MONGIN (1995): "A Note on Affine Aggregation," *Economics Letters*, 47, 177–183.
- DEKEL, E., B. L. LIPMAN, AND A. RUSTICHINI (2001): "Representing Preferences with a Unique Subjective State Space," *Econometrica*, 69, 891–934.
- DEKEL, E., B. L. LIPMAN, A. RUSTICHINI, AND T. SARVER (2007): "Representing Preference with a Unique Subjective State Space: Corrigendum," *Econometrica*, 75, 591–600.
- FISHBURN, P. (1978): "Axioms for Approval Voting: Direct Proof," *Journal of Economic Theory*, 19, 180–185.
- GUL, F., AND W. PESENDORFER (2001): "Temptation and Self-Control," *Econometrica*, 69, 1403–1435.
- HARSANYI, J. C. (1955): "Cardinal Welfare, Individualistic Ethics, and Interpersonal Comparisons of Utility," *Journal of Political Economy*, 63, 309–321.
- KREPS, D. M. (1979): "A Representation Theorem for 'Preference for Flexibility'," *Econometrica*, 47, 565–576.

WEBER, R. J. (1995): "Approval Voting," *Journal of Economic Perspectives*, 9, 39-49.